

Failure of mean field approximation in weakly coupled dilaton gravity*

Fitkevich M. D.

Institute for Nuclear Research of RAS
Moscow Institute of Physics and Technology




Quarks–2018
May 29, Valday

* - *in collaboration with D.G. Levkov, S.M. Sibiryakov, and Y.A. Zenkevich*

Motivation

- Information paradox: apparent loss of quantum coherence.



$$|out\rangle = \hat{S}|in\rangle, \quad \hat{S}\hat{S}^\dagger \neq \hat{1}$$

Hawking, 1975

- ▶ Responses: AdS/CFT correspondence, “complementarity”, etc.
- ▶ **AMPS-firewall**. Unitarity versus Equivalence principle.

Almheiri et al, 2012

We need useful solvable models!

- Toy models: 2D dilaton gravity. Period of activity: $t \in (1991, 1996)$.
Problem: apparent non-unitarity persists.
- Idea: revive 2d gravity with new semi-classical methods!
 - ▶ S-matrix as path integral: complex classical solutions.

Weakly coupled dilaton gravity

$$S = \int d^2x \sqrt{-g} \left[e^{-2\phi} \left(R + 4(\nabla\phi)^2 + 4\lambda^2 \right) - \frac{1}{2}(\nabla f)^2 \right]$$


ArXiv:9111056 [hep-th] C. Callan, S. Giddings, J. Harvey, A. Strominger, 1991

In the bulk:

$$ds^2 = -e^{2\phi} dvdu, \quad f(v, u) = f_{out}(u) + f_{in}(v)$$

$$e^{-2\phi} = -\lambda^2 vu - \mathcal{T}(v) - \mathcal{H}(u)$$

$$\partial_v^2 \mathcal{T} = (\partial_v f_{in})^2/2, \quad \partial_u^2 \mathcal{H} = (\partial_u f_{out})^2/2$$



Weakly coupled dilaton gravity

$$S = \int d^2x \sqrt{-g} \left[e^{-2\phi} \left(R + 4(\nabla\phi)^2 + 4\lambda^2 \right) - \frac{1}{2}(\nabla f)^2 \right] + 2 \int_{\partial\mathcal{M}} d\tau e^{-2\phi} (K + 2\lambda)$$

ArXiv:9111056 [hep-th] C. Callan, S. Giddings, J. Harvey, A. Strominger, 1991

ArXiv:1702.02576 [hep-th] M.F., D. Levkov, Y. Zenkevich, 2017

In the bulk:

$$ds^2 = -e^{2\phi} dvdu, \quad f(v, u) = f_{out}(u) + f_{in}(v)$$

$$e^{-2\phi} = -\lambda^2 vu - \mathcal{T}(v) - \mathcal{H}(u)$$

$$\partial_v^2 \mathcal{T} = (\partial_v f_{in})^2 / 2, \quad \partial_u^2 \mathcal{H} = (\partial_u f_{out})^2 / 2$$


- Weak coupling: $g_{gr} = e^\phi \leq e^{\phi_0} \ll 1$

- Minimal black hole mass


$$M_{cr} = 2\lambda e^{-2\phi_0}$$

- Reflecting condition

$$f_{out}(U(v)) = f_{in}(v).$$



One-loop effective action




Includes Hawking radiation and backreaction on metric.


One loop from $N = 24Q^2$ scalars $f_i \Rightarrow$
Liouville-Polyakov action

$$S_{PL} = -\frac{Q^2}{2} \int dx \sqrt{-g} \int dy \sqrt{-g} R \frac{1}{\square} R$$

$$S_{PL} \underset{\text{on-shell}}{\equiv} \int d^2x \sqrt{-g} \left[-\frac{1}{2} (\nabla \chi)^2 + Q \chi R \right]$$



One-loop effective action



Includes Hawking radiation and backreaction on metric.

One loop from $N = 24Q^2$ scalars $f_i \Rightarrow$ Liouville-Polyakov action

$$S_{PL} = -\frac{Q^2}{2} \int dx \sqrt{-g} \int dy \sqrt{-g} R \frac{1}{\square} R$$

$$S_{PL} \underset{\text{on-shell}}{\equiv} \int d^2x \sqrt{-g} \left[-\frac{1}{2} (\nabla \chi)^2 + Q \chi R \right]$$


To restore solvability: *ArXiv:9206070 [hep-th]*

J. Russo, L. Susskind, L. Thorlacius, 1992

$$\Delta S_{RST} = -Q^2 \int d^2x \sqrt{-g} \phi R$$

Boundary terms: fixed by **Wess-Zumino**

$$\Delta S_{\text{boundary}} = 2 \int d\tau \left[(-Q^2 \phi + Q \chi) K + \lambda Q^2 \right]$$




M.F., D. Levkov, Y. Zenkevich, in preparation

Solution for black hole

$U(v)$ - boundary in light-cone coordinates u, v ; $T(v)$ - incoming matter f_{in} .

$$\text{Boundary condition: } \partial_v U = \text{const} \left(\partial_v T + \lambda^2 U + \frac{Q^2 \partial_v^2 U}{2\partial_v U} \right)^2$$

$E > E_{cr} \Rightarrow$ 2nd branch $\bar{U}(v) \neq U(v)$ of $\phi(v, u) = \phi_0$ is the singularity.




Solution for black hole

$U(v)$ - boundary in light-cone coordinates u, v ; $T(v)$ - incoming matter f_{in} .

$$\text{Boundary condition: } \partial_v U = \text{const} \left(\partial_v T + \lambda^2 U + \frac{Q^2 \partial_v^2 U}{2\partial_v U} \right)^2$$

$E > E_{cr} \Rightarrow$ 2nd branch $\bar{U}(v) \neq U(v)$ of $\phi(v, u) = \phi_0$ is the singularity.

Impose new boundary condition \Rightarrow new boundary.



Solution for black hole

$U(v)$ - boundary in light-cone coordinates u, v ; $T(v)$ - incoming matter f_{in} .

$$\text{Boundary condition: } \partial_v U = \text{const} \left(\partial_v T + \lambda^2 U + \frac{Q^2 \partial_v^2 U}{2\partial_v U} \right)^2$$


$E > E_{cr} \Rightarrow$ 2nd branch $\bar{U}(v) \neq U(v)$ of $\phi(v, u) = \phi_0$ is the singularity.

Impose new boundary condition \Rightarrow new boundary.

Problems:


- Non-analyticity.
- Ambiguity: $\partial_v U \Big|_{\text{end point}} - ?$
- Thunderpop: $E_{th} \sim -\lambda Q^2$.

\Rightarrow has to introduce smearing around thunderpop.



Failure of mean field theory

Vacuum correlation function $G_{vac}(\bar{v}, \bar{v}') \equiv \langle f(\bar{v})f(\bar{v}') \rangle \propto \ln |\bar{v} - \bar{v}'|$




Near the thunderpop $\bar{u}, \bar{u}' \simeq \bar{u}_{end}$,
 $G(\bar{u}, \bar{u}') = \langle f_{out}(\bar{u})f_{in}(\bar{u}') \rangle = \langle f_{in}(\bar{v}(\bar{u}))f_{in}(\bar{v}(\bar{u}')) \rangle \neq G_{vac}(\bar{u}, \bar{u}')$

\Rightarrow thunderbolt: particles with **arbitrary large momenta** $k \sim \Delta\bar{u}^{-1}$.

Strominger, 1994

Energy conservation $\Rightarrow \Delta\bar{u} \simeq Q/M_{cr}$ - characteristic size of quantum area where semiclassicals always fails.

Failure of mean field theory



$$\langle \Psi_{out} | \hat{S} | \Psi_{in} \rangle = \int \mathcal{D}\Phi \Psi_{out}^* \Psi_{in} \exp\left\{\frac{i}{\hbar} S[\Phi]\right\}, \quad \Phi = \{g_{\mu\nu}, \phi, f\}$$

Semiclassics $\Rightarrow \frac{\delta}{\delta\Phi} S = 0 \Rightarrow$ saddle point Φ_s - can not be singular.

- Mean field approximation fails: glued solutions are **incorrect saddle points!**
- Another possible answer: stiff boundary condition is inconsistent.
 - ▶ Analogy: Klein paradox in QM. Second quantization of the boundary?

Calculating S-matrix elements

- But we want to consider the whole solution Φ_s corresponding to $\Psi_{in} \mapsto \Psi_{out}$ (asymptotically flat to asymptotically flat).
 - ▶ At $E < E_{cr} \Rightarrow$ semiclassical S-matrix exists: $\langle \Psi_{out} | \hat{S} | \Psi_{in} \rangle \approx \exp\{\frac{i}{\hbar} S[\Phi_c]\}$
 - ▶ At $E > E_{cr} \Rightarrow$ no flat asymptotics of classical solutions at $t \rightarrow +\infty \Rightarrow$ ill-defined S-matrix.
- Idea: obtain physical solutions at $E > E_{cr}$ via analytic continuation.
 - ▶ Problem: many possibilities for deformation: suppressed, unphysical solutions. We need leading contribution!

We need a criterion to chose physical branch at $E > E_{cr}$, $\Im m E \rightarrow 0$.

Criterion: start from the “shell” model


$$S = -m \int d\tau \sqrt{-g_{\mu\nu} \frac{dy^\mu}{d\tau} \frac{dy^\nu}{d\tau}} + S_{gravity}$$

Junction condition on shell: $\left(\frac{dr}{d\tau}\right)^2 - \left(\frac{M}{m} + \frac{m}{8\lambda} e^{-2\lambda r}\right)^2 + 1 = 0$

Well-known analytic continuation: $M \mapsto M + i\varepsilon, \quad \varepsilon \rightarrow +0$

ArXiv:9907001 [hep-th] M. Parikh, F. Wilczek, 2000

ArXiv:1503.07181 [hep-th] F. Bezrukov, D. Levkov, S. Sibiriakov, 2015




Can be generalized to field theory.

“Shell” model: result

$$2\Im m S_{tot} = S_{BH} - S_{cr}, \quad S = \frac{M}{T_H} = \frac{2\pi}{\lambda} M$$

Plots for $m = 0$:



Probability:

$$\mathcal{P}_{fi} \approx \exp(-S_{BH} + S_{cr}).$$

Non-entropic suppression: Nature abhors discontinuity.

Conclusions

- Mean field theory:
 - ▶ Either not a good approximation...
 - ▶ ...or models with stiff boundaries are not self-consistent.
- Complex semiclassical method:
 - ▶ Reliable analytic continuation for shells.
 - ▶ Relevant solutions for fields.
 - ▶ Non-entropic suppression.

Thank you for attention!