

ATLAS status & overview



Evgenii Baldin

Budker Institute of Nuclear Physics,
Novosibirsk State University

for the ATLAS Collaboration
Quarks-2018 (27 May – 2 June, 2018)



- 1 Table of Contents
- 2 LHC status and plan
- 3 ATLAS detector
- 4 Higgs Results
- 5 Standard Model Results
- 6 Top Physics Results
- 7 Heavy Ions Results
- 8 Supersymmetry Search
- 9 Exotics Search
- 10 Conclusion

LHC / HL-LHC Plan



- Most of **Run 1** (7 and 8 TeV) analyzes were done.



LHC / HL-LHC Plan



integrated luminosity


- Most of **Run 1** (7 and 8 TeV) analyzes were done.
- Newest results mostly related to 2015-2016 statistics (**Run 2**, 36 fb⁻¹)

LHC / HL-LHC Plan




- Most of **Run 1** (7 and 8 TeV) analyzes were done.
- Newest results mostly related to 2015-2016 statistics (**Run 2**, 36 fb^{-1})
- This year data taking campaign started successfully (**Run 2** 150 fb^{-1} is achievable by the end of 2018)



LHC / HL-LHC Plan




- Most of **Run 1** (7 and 8 TeV) analyzes were done.
- Newest results mostly related to 2015-2016 statistics (**Run 2**, 36 fb⁻¹)
- This year data taking campaign started successfully (**Run 2** 150 fb⁻¹ is achievable by the end of 2018)
- Probably full **Run 2** (13 TeV) statistics analyzes will be ready just when **Run 3** (14 TeV) will be started.

ATLAS detector






Detector performance with pile-up





40 $\langle\mu\rangle$ corresponds to 20 primary vertices



Electron identification efficiencies

The situation with pile-up looks adequate even for **Run 3**.


Measurement of the Higgs boson mass $\sqrt{s} = 13$ TeV



$\sqrt{s} = 13$ TeV, 36.1 fb^{-1}

ATLAS-CONF-2017-046


Measurement of the Higgs boson mass $\sqrt{s} = 13$ TeV



$\sqrt{s} = 13$ TeV, 36.1 fb^{-1}

ATLAS-CONF-2017-046



Higgs boson production



Total $pp \rightarrow H + X$ cross section as function of \sqrt{s} compared to Standard Model prediction at up to N³LO in QCD.

ATLAS-CONF-2017-047

Couplings in $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^{(*)} \rightarrow 4\ell$





Cross sections for ggF, VBF, VH, and $t\bar{t}H$ normalized to the SM predictions and measured with the assumption of SM branching fractions. Overall good agreement except for the VBF.

Quantity Results of a simultaneous fit for $\sigma_{ggF} \cdot B_{4\ell}$, $B_{\gamma\gamma}/B_{4\ell}$, $\sigma_{VBF}/\sigma_{ggF}$, σ_{VH}/σ_{ggF} , and $\sigma_{t\bar{t}H}/\sigma_{ggF}$. The fit results are normalized to the SM predictions.

ATLAS-CONF-2017-047

Evidence for the $H \rightarrow b\bar{b}$ decay with the ATLAS detector





Observed significance: 3.5σ (exp. 3.0σ).

The fitted values of the Higgs boson signal strength parameter μ for $m_H = 125 \text{ GeV}$ for the 0-, 1- and 2-lepton channels and their combination.

$$\sqrt{s} = 13 \text{ TeV}, \mathcal{L} = 36.1 \text{ fb}^{-1}.$$

arXiv:1708.03299

Evidence for the $H \rightarrow b\bar{b}$ decay with the ATLAS detector



Observed significance: 3.6σ (exp. 4.0σ).


Event yields as a function of $\log(S/B)$ for data, background (B) and Higgs boson signal (S) with $m_H = 125 \text{ GeV}$, for the 7 TeV, 8 TeV and 13 TeV datasets combined. Multivariate analysis based on event kinematics was used.

arXiv:1708.03299

Tests of the standard model by ATLAS

Standard Model Production Cross Section Measurements

Status: March 2014




March 2014

Tests of the standard model by ATLAS

Standard Model Production Cross Section Measurements

Status: March 2018




March 2018

Tests of the standard model by ATLAS (Diboson)

Diboson Cross Section Measurements


Status: July 2017



	$\int \mathcal{L} dt$ [fb $^{-1}$]	Reference
20.2	PRD 95 (2017) 112005	
4.9	JHEP 01, 086 (2013)	
4.6	PRD 87, 112003 (2013)	
4.6	arXiv:1407.1618 [hep-ph]	
4.6	PRD 87, 112003 (2013)	
20.3	PRD 93, 112002 (2016)	
4.6	PRD 87, 112002 (2013)	
4.6	arXiv:1407.1618 [hep-ph]	
20.3	PRD 93, 112002 (2016)	
4.6	PRD 87, 112003 (2013)	
20.3	PRD 93, 112002 (2016)	
4.6	PRD 87, 112003 (2013)	
20.2	arXiv: 1706.01702 [hep-ex]	
4.6	JHEP 01, 049 (2015)	
20.2	arXiv: 1706.01702 [hep-ex]	
3.2	arXiv: 1702.04519 [hep-ex]	
20.3	PLB 763, 114 (2016)	
4.6	PRD 87, 112001 (2013)	
4.6	PRL 113, 212001 (2014)	
3.2	arXiv: 1702.04519 [hep-ex]	
20.3	JHEP 09 (2016) 029	
4.6	PRD 87, 112001 (2013)	
4.6	PRD 91, 052005 (2015)	
20.3	PLB 763, 114 (2016)	
3.2	PLB 762 (2016) 1	
20.3	PLB 761 (2016) 179	
4.6	EPLC 72, 2173 (2012)	
3.2	PLB 762 (2016) 1	
20.3	PRD 93, 092004 (2016)	
36.1	ATLAS-CONF-2017-031	
20.3	PLB 735 (2014) 311	
20.3	JHEP 01, 098 (2017)	
4.6	JHEP 01, 028 (2013)	
36.1	ATLAS-CONF-2017-031	
20.3	JHEP 01, 099 (2017)	
4.6	JHEP 03, 128 (2013)	
20.3	JHEP 01, 099 (2017)	
4.6	JHEP 03, 128 (2013)	
20.3	PLB 753, 552-572 (2016)	
4.6	JHEP 03, 128 (2013)	

Data prefers NNLO over NLO!

SM cross-section measurements



Summary of total production cross-section measurements by ATLAS presented as a function of centre-of-mass energy \sqrt{s} from 7 to 13 TeV for a few selected processes.

Measurement of the W boson mass

Collected at CERN LHC with ATLAS detector in 2011, $\sqrt{s} = 7 \text{ TeV}$

$W \rightarrow \mu\nu$ candidates: $7.8 \cdot 10^6$

$W \rightarrow e\nu$ candidates: $5.9 \cdot 10^6$

The mass of the W boson can be expressed in terms of the other SM parameters as follows:

$$m_W^2 \left(1 - \frac{m_W^2}{m_Z^2}\right) = \frac{\pi\alpha}{\sqrt{2}G_\mu} (1 + \Delta r),$$


where Δr incorporates effect of high order corrections.

PDG average $m_W = 80385 \pm 15 \text{ MeV}$ (mainly CDF and D0)

SM prediction $m_W = 80356 \pm 8 \text{ MeV}$ (based on arXiv:1407.3792 with updated m_t and m_H)

arXiv:1701.07240

Measurement of the W boson mass




$$m_W = 80370 \pm 7(\text{stat}) \pm 11(\text{syst}) \pm 14(\text{mod}) \text{ MeV} = 80370 \pm 19 \text{ MeV}$$

$$m_{W^+} - m_{W^-} = -29 \pm 12.8(\text{stat}) \pm 7.0(\text{syst}) \pm 23.9(\text{mod}) \text{ MeV} = -29 \pm 28 \text{ MeV}$$

arXiv:1701.07240

Measurement of the W boson mass




$$m_W = 80370 \pm 7(\text{stat}) \pm 11(\text{syst}) \pm 14(\text{mod}) \text{ MeV} = 80370 \pm 19 \text{ MeV}$$

$$m_{W^+} - m_{W^-} = -29 \pm 12.8(\text{stat}) \pm 7.0(\text{syst}) \pm 23.9(\text{mod}) \text{ MeV} = -29 \pm 28 \text{ MeV}$$


arXiv:1701.07240

Top-pair production cross-section




Summary of LHC and Tevatron measurements of the top-pair production cross-section as a function of the centre-of-mass energy compared to the NNLO QCD calculation complemented with NNLL resummation.

Single top production cross-section



Summary of ATLAS measurements of the single top production cross-sections in the t-channel as a function of the center of mass energy compared to a theoretical calculation based on NLO QCD.



Summary of the ATLAS direct m_{top} measurements



The results are compared to the ATLAS, Tevatron and Tevatron+LHC m_{top} combinations.

ATLAS-CONF-2017-071

Associated top and Z production (first observation)



WWZ and tZ couplings probe


Post-fit neural-network output distributions in the signal region.

$$\sigma_{tZ}^{\text{ATLAS}} = 600 \pm 170 \text{ (stat.)} \pm 140 \text{ (syst.)} \text{ fb.} \quad (\text{Observed, } 4.2\sigma)$$


$$\sigma_{tZ}^{\text{SM}} = 800^{+49}_{-59} \text{ fb.} \quad (\text{Expected significance } 5.4\sigma)$$

arXiv:1710.03659

Evidence for light-by-light scattering in heavy-ion collisions



Ultra-peripheral heavy ion collisions




$$\sigma_{\text{fid}}^{\text{ATLAS}} = 70 \pm 24 \text{ (stat.)} \pm 17 \text{ (syst.) nb},$$


$$\sigma_{\text{fid}}^{\text{SM}} = 45 \pm 9 \text{ nb}.$$

arXiv:1702.01625

J/ψ and $\psi(2S)$ suppression in 5.02 TeV Pb+Pb collisions





Comparison of prompt $J/\psi R_{AA}$ (nuclear modification factor) measured in 5.02 TeV PbPb collisions by ATLAS with the inclusive $J/\psi R_{AA}$ measured by ALICE.



$\psi(2S)$ to J/ψ double ratio, as a function of the number of participants
The suppression of prompt $\psi(2S)$ is stronger than that of J/ψ .

arXiv:1805.04077

squarks and gluinos




(squark, lightest neutralino) mass plane

(gluino, lightest neutralino) mass plane

Exclusion limits at 95% CL in the mass plane for different simplified models.

Higgsinos search

March 2018



Exclusion limits at 95% CL for higgsino pair production $\chi_1^+ \chi_1^-$, $\chi_1^\pm \chi_1^0$, $\chi_1^1 \chi_2^0$, and $\chi_1^0 \chi_2^0$ with off-shell SM-boson-mediated decays to the lightest neutralino, χ_1^0 , as a function of the χ_1^\pm and χ_1^0 masses. The production cross-section is for pure higgsinos.

arXiv:1712.08119

ATLAS Exotics Searches

ATLAS Exotics Searches* - 95% CL Upper Exclusion Limits


Status: July 2017

ATLAS Preliminary

$$\int \mathcal{L} dt = (3.2 - 37.0) \text{ fb}^{-1}$$

$$\sqrt{s} = 8, 13 \text{ TeV}$$


Reference



*Only a selection of the available mass limits on new states or phenomena is shown.


†Small-radius (large-radius) jets are denoted by the letter j (J).

Dark Matter searches




Regions in a dark matter mass-mediator mass plane excluded at 95% CL by a selection of ATLAS dark matter searches, for one possible interaction between the Standard Model and dark matter, the leptophobic axial-vector mediator as described in arXiv:1703.05703.

A search for lepton-flavor-violating Z decays



1-prong $\tau_{\text{had-vis}}$ candidates



3-prong $\tau_{\text{had-vis}}$ candidates

Neural network classifiers are used to discriminate signal from backgrounds, and the NN output distributions are analyzed in a template fit to data.

$$\mathcal{B}(Z \rightarrow e\tau) < 5.8 \cdot 10^{-5} \text{ and } \mathcal{B}(Z \rightarrow \mu\tau) < 2.4 \cdot 10^{-5}.$$

arXiv:1804.09568

- LHC and ATLAS are in a good shape. **Run 2** will end with the end of 2018. At least for now there are no problems preventing the implementation of the plan for the data taking (150 fb^{-1}).
- Standard Model has been probed with unprecedented precision. No deviation from the Standard Model prediction is observed so far.
- Please see “ATLAS experiment — public results” web page for more information.

▶ [Link to "ATLAS experiment — public results"](#)