


Recent developments in charmed baryon spectroscopy

Elena Solovieva

Moscow Institute of Physics and Technology
Lebedev Physical Institute of the RAS

Known Charmed Baryon States



$$\mathcal{B}_c = c + \text{diquark}$$

Quark content of diquark:

- qq with isospin 0 (flavor antisymmetric) — **Λ_c family**;
- qq with isospin 1 (flavor symmetric) — **Σ_c family**;
- qs with isospin $\frac{1}{2}$ — **Ξ_c family**;
- ss with isospin 0 (flavor symmetric) — **Ω_c family**.



Ξ_c Family: PDG'2016

State	Decay mode	Mass, MeV/ c^2	Width, MeV	J^P
Ξ_c' ⁺	$\Xi_c^+ \gamma$	2575.7 ± 3.0		$\frac{1}{2}^+$
Ξ_c' ⁰	$\Xi_c^0 \gamma$	2577.9 ± 2.9		$\frac{1}{2}^+$
$\Xi_c(2645)^+$	$\Xi_c^0 \pi^+$	2645.9 ± 0.5	2.6 ± 0.4	$\frac{3}{2}^+$
$\Xi_c(2645)^0$	$\Xi_c^+ \pi^-$	2645.9 ± 0.5	$< 5.5 @ 90\% \text{ CL}$	$\frac{3}{2}^+$
$\Xi_c(2790)^+$	$\Xi_c'^0 \pi^+$	2789.1 ± 3.2	$< 15 @ 90\% \text{ CL}$	$\frac{1}{2}^-$
$\Xi_c(2790)^0$	$\Xi_c'^+ \pi^-$	2791.9 ± 3.3	$< 12 @ 90\% \text{ CL}$	$\frac{1}{2}^-$
$\Xi_c(2815)^+$	$\Xi_c^+ \pi^+ \pi^-$, $\Xi_c(2645)^0 \pi^+$	2816.6 ± 0.9	$< 3.5 @ 90\% \text{ CL}$	$\frac{3}{2}^-$
$\Xi_c(2815)^0$	$\Xi_c^0 \pi^+ \pi^-$, $\Xi_c(2645)^+ \pi^-$	2819.6 ± 1.2	$< 6.5 @ 90\% \text{ CL}$	$\frac{3}{2}^-$
$\Xi_c(2930)^0$	$\Lambda_c^+ K^-$	2931 ± 6	36 ± 13	
$\Xi_c(2970)^+$	$\Lambda_c^+ K^- \pi^+$, $\Sigma_c^{++} K^-$, $\Xi_c(2645)^0 \pi^+$	2970.7 ± 2.2	17.9 ± 3.5	
$\Xi_c(2970)^0$	$\Xi_c(2645)^+ \pi^-$	2968.0 ± 2.6	20 ± 7	
$\Xi_c(3055)^+$	$\Sigma_c^{++} K^-$	3055.1 ± 1.7	11 ± 4	
$\Xi_c(3080)^+$	$\Lambda_c^+ K^- \pi^+$, $\Sigma_c^{++} K^-$, $\Sigma_c(2520)^{++} K^-$	3076.94 ± 0.28	4.3 ± 1.5	
$\Xi_c(3080)^0$	$\Lambda_c^+ K_S^0 \pi^-$, $\Sigma_c^0 K_S^0$, $\Sigma_c(2520)^0 K_S^0$	3079.9 ± 1.4	5.6 ± 2.2	#3

Decays to Ξ_c' : Ξ_c' Isodoublet

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$

$$\Xi_c' \rightarrow \Xi_c \gamma$$





$$M_{\Xi_c'} - M_{\Xi_c^+} = (110.5 \pm 0.1[\text{stat.}] \pm 0.4[\text{syst.}]) \text{ MeV}/c^2$$

$$M_{\Xi_c'} - M_{\Xi_c^0} = (108.3 \pm 0.1[\text{stat.}] \pm 0.4[\text{syst.}]) \text{ MeV}/c^2$$

Decays to Ξ_c : $\Xi_c(2790)$ Isodoublet

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$

$$\Xi_c(2790) \rightarrow \Xi_c' \pi$$



$$M_{\Xi_c(2790)^+} - M_{\Xi_c'^0} = (213.2 \pm 0.2[\text{stat.}] \pm 0.1[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2790)^+} = (8.9 \pm 0.6[\text{stat.}] \pm 0.8[\text{syst.}]) \text{ MeV}$$

$$M_{\Xi_c(2790)^0} - M_{\Xi_c'^+} = (215.7 \pm 0.2[\text{stat.}] \pm 0.1[\text{syst.}]) \text{ MeV}/c^2$$



$$\Gamma_{\Xi_c(2790)^0} = (10.0 \pm 0.7[\text{stat.}] \pm 0.8[\text{syst.}]) \text{ MeV}$$

Decays to Ξ_c : $\Xi_c(2645)$ Isodoublet

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$

$$\Xi_c(2645) \equiv \Xi_c^*$$

$$\Xi_c(2645) \rightarrow \Xi_c \pi$$



$$M_{\Xi_c(2645)^+} - M_{\Xi_c^0} = (174.66 \pm 0.06[\text{stat.}] \pm 0.07[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2645)^+} = (2.06 \pm 0.13[\text{stat.}] \pm 0.13[\text{syst.}]) \text{ MeV}$$



$$M_{\Xi_c(2645)^0} - M_{\Xi_c^+} = (178.46 \pm 0.07[\text{stat.}] \pm 0.07[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2645)^0} = (2.35 \pm 0.18[\text{stat.}] \pm 0.13[\text{syst.}]) \text{ MeV}$$

Decays to Ξ_c : $\Xi_c(2815)$ Isodoublet

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$

$$\Xi_c(2815) \rightarrow \Xi_c^* \pi$$



$$M_{\Xi_c(2815)^+} - M_{\Xi_c^+} = (348.80 \pm 0.08[\text{stat.}] \pm 0.06[\text{syst.}]) \text{ MeV}/c^2$$




$$\Gamma_{\Xi_c(2815)^+} = (2.43 \pm 0.20[\text{stat.}] \pm 0.17[\text{syst.}]) \text{ MeV}$$

$$M_{\Xi_c(2815)^0} - M_{\Xi_c^0} = (349.35 \pm 0.08[\text{stat.}] \pm 0.07[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2815)^0} = (2.54 \pm 0.18[\text{stat.}] \pm 0.17[\text{syst.}]) \text{ MeV}$$

Decays to Ξ_c : $\Xi_c(2970)$ Isodoublet

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$



$$M_{\Xi_c(2970)^+} - M_{\Xi_c^+} = (498.1 \pm 0.8[\text{stat.}] \pm 0.2[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2970)^+} = \left(28.1 \pm 2.4[\text{stat.}] \begin{array}{l} +1.0 \\ -5.0 \end{array} [\text{syst.}] \right) \text{ MeV}$$

$$M_{\Xi_c(2970)^0} - M_{\Xi_c^0} = (499.9 \pm 0.7[\text{stat.}] \pm 0.2[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2970)^0} = \left(30.3 \pm 2.3[\text{stat.}] \begin{array}{l} +1.0 \\ -1.8 \end{array} [\text{syst.}] \right) \text{ MeV}$$

Ξ_c Family: Decays to Ξ_c

Particle	Yield	Mass	$M - M(\Xi_c)$	$M - M(\Xi'_c)$	Width
$\Xi_c^{'+}$ PDG	7055 \pm 211	$2578.4 \pm 0.1 \pm 0.4^{+0.3}_{-0.4}$	$110.5 \pm 0.1 \pm 0.4$		
		2575.6 ± 3.0	107.8 ± 3.0		
Ξ_c^0 PDG	11560 \pm 276	$2579.2 \pm 0.1 \pm 0.4^{+0.3}_{-0.4}$	$108.3 \pm 0.1 \pm 0.4$		
		2577.9 ± 2.9	107.0 ± 2.9		
$\Xi_c(2645)^+$ PDG	1260 \pm 40	$2645.58 \pm 0.06 \pm 0.07^{+0.28}_{-0.40}$	$174.66 \pm 0.06 \pm 0.07$		$2.06 \pm 0.13 \pm 0.13$
		2645.9 ± 0.5	175.0 ± 0.6		$2.6 \pm 0.2 \pm 0.4$
$\Xi_c(2645)^0$ PDG	975 \pm 36	$2646.43 \pm 0.07 \pm 0.07^{+0.28}_{-0.40}$	$178.46 \pm 0.07 \pm 0.07$		$2.35 \pm 0.18 \pm 0.13$
		2645.9 ± 0.5	178.0 ± 0.6		< 5.5
$\Xi_c(2790)^+$ PDG	2231 \pm 103	$2791.6 \pm 0.2 \pm 0.1 \pm 0.4^{+0.3}_{-0.4}$	$320.7 \pm 0.2 \pm 0.1 \pm 0.4$	$213.2 \pm 0.2 \pm 0.1$	$8.9 \pm 0.6 \pm 0.8$
		2789.8 ± 3.2	318.2 ± 3.2		< 15
$\Xi_c(2790)^0$ PDG	1241 \pm 72	$2794.9 \pm 0.3 \pm 0.1 \pm 0.4^{+0.3}_{-0.4}$	$323.8 \pm 0.2 \pm 0.1 \pm 0.4$	$215.7 \pm 0.2 \pm 0.1$	$10.0 \pm 0.7 \pm 0.8$
		2791.9 ± 3.3	324.0 ± 3.3		< 12
$\Xi_c(2815)^+$ PDG	941 \pm 35	$2816.73 \pm 0.08 \pm 0.06^{+0.28}_{-0.40}$	$348.80 \pm 0.08 \pm 0.06$		$2.43 \pm 0.20 \pm 0.17$
		2816.6 ± 0.9	348.7 ± 0.9		< 3.5
$\Xi_c(2815)^0$ PDG	1258 \pm 40	$2820.20 \pm 0.08 \pm 0.07^{+0.28}_{-0.40}$	$349.35 \pm 0.08 \pm 0.07$		$2.54 \pm 0.18 \pm 0.17$
		2819.6 ± 1.2	348.8 ± 1.2		< 6.5
$\Xi_c(2970)^+$ PDG	916 \pm 55	$2966.0 \pm 0.8 \pm 0.2^{+0.3}_{-0.4}$	$498.1 \pm 0.8 \pm 0.2$		$28.1 \pm 2.4^{+1.0}_{-5.0}$
		2970.7 ± 2.2			17.9 ± 3.5
$\Xi_c(2970)^0$ PDG	1443 \pm 75	$2970.8 \pm 0.7 \pm 0.2^{+0.3}_{-0.4}$	$499.9 \pm 0.7 \pm 0.2$		$30.3 \pm 2.3^{+1.0}_{-1.8}$
		$2968.0 \pm 2.6 \pm 0.5$			20 ± 7

[J. Yelton *et al.* (Belle Collaboration), Phys. Rev. D **94**, 052011 (2016)]


Ξ_c Family: Decays to $\Lambda_c(\Sigma_c)$

Ξ_c^+	
Ξ_c^0	
$\Xi_c(2645)^+$	
$\Xi_c(2645)^0$	
$\Xi_c(2790)^+$	
$\Xi_c(2790)^0$	
$\Xi_c(2815)^+$	
$\Xi_c(2815)^0$	
$\Xi_c(2930)^0$	
$\Xi_c(2970)^+$	
$\Xi_c(2970)^0$	
$\Xi_c(3055)^+$	
$\Xi_c(3055)^0$	
$\Xi_c(3080)^+$	
$\Xi_c(3080)^0$	


$$B^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^- K^-$$

$$m_{\Xi_c(2930)^0} = (2931 \pm 3[\text{stat.}] \pm 5[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2930)^0} = (36 \pm 7[\text{stat.}] \pm 11[\text{syst.}]) \text{ MeV}$$



[B. Aubert *et al.* (BaBar Collaboration),
Phys. Rev. D **77**, 031101 (2008)]





$$m_{\Xi_c(2930)^0} = (2928.9 \pm 3.0[\text{stat.}]^{+0.9}_{-12.0}[\text{syst.}]) \text{ MeV}/c^2$$

$$\Gamma_{\Xi_c(2930)^0} = (19.5 \pm 8.4[\text{stat.}]^{+5.9}_{-7.9}[\text{syst.}]) \text{ MeV}$$

[Y.B. Li, C.P. Shen *et al.* (Belle Collaboration),
Eur. Phys. J. C **78**, 252 (2018)]

Ξ_c Family: Decays to $\Lambda_c(\Sigma_c)$



Ξ_c^+	Ξ_c^0
$\Xi_c(2645)^+$	
$\Xi_c(2645)^0$	
$\Xi_c(2790)^+$	
$\Xi_c(2790)^0$	
$\Xi_c(2815)^+$	
$\Xi_c(2815)^0$	
$\Xi_c(2930)^0$	
$\Xi_c(2970)^+$	
$\Xi_c(2970)^0$	
$\Xi_c(3055)^+$	
$\Xi_c(3055)^0$	
$\Xi_c(3080)^+$	
$\Xi_c(3080)^0$	



Particle	Mass (MeV/c ²)	Width (MeV/c ²)
$\Xi_c(2970)^+$	$2974.9 \pm 1.5 \pm 2.1$	$14.8 \pm 2.5 \pm 4.1$
$\Xi_c(3055)^+$	$3058.1 \pm 1.0 \pm 2.1$	$9.7 \pm 3.4 \pm 3.3$
$\Xi_c(3080)^+(\Sigma_c)$	$3077.9 \pm 0.4 \pm 0.7$	$3.2 \pm 1.3 \pm 1.3$
$\Xi_c(3080)^+(\Sigma_c^*)$	$3076.9 \pm 0.3 \pm 0.2$	$2.4 \pm 0.9 \pm 1.6$

Ξ_c Family: Decays to $\Lambda_c(\Sigma_c)$

Ξ_c^+
Ξ_c^0
$\Xi_c(2645)^+$
$\Xi_c(2645)^0$
$\Xi_c(2790)^+$
$\Xi_c(2790)^0$
$\Xi_c(2815)^+$
$\Xi_c(2815)^0$
$\Xi_c(2930)^0$
$\Xi_c(2970)^+$
$\Xi_c(2970)^0$
$\Xi_c(3055)^+$
$\Xi_c(3055)^0$
$\Xi_c(3080)^+$
$\Xi_c(3080)^0$







[Y. Kato, T. Iijima *et al.* (Belle Collaboration), Phys. Rev. D **89**, 052003 (2014)]

[B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **77**, 012002 (2008)]

Ξ_c Family: Decays to ΛD

Ξ_c^+	Ξ_c^0
$\Xi_c(2645)^+$	$\Xi_c(2645)^0$
$\Xi_c(2790)^+$	$\Xi_c(2790)^0$
$\Xi_c(2815)^+$	$\Xi_c(2815)^0$
$\Xi_c(2930)^0$	
$\Xi_c(2970)^+$	
$\Xi_c(2970)^0$	
$\Xi_c(3055)^+$	
$\Xi_c(3055)^0$	
$\Xi_c(3080)^+$	
$\Xi_c(3080)^0$	


Resonance	Mass (MeV/c^2)	Width (MeV)	Significance (σ)
$\Xi_c(3055)^0$	$3059.0 \pm 0.5 \pm 0.6$	$6.4 \pm 2.1 \pm 1.1$	8.6
$\Xi_c(3055)^+$	$3055.8 \pm 0.4 \pm 0.2$	$7.0 \pm 1.2 \pm 1.5$	11.7
$\Xi_c(3080)^+$	$3079.6 \pm 0.4 \pm 0.1$	< 6.3	4.8



Ξ_c Family

State	Decay mode	Mass, MeV/ c^2	Width, MeV	J^P
$\Xi_c'^+$	$\Xi_c^+ \gamma$	2577.4 ± 1.2		$\frac{1}{2}^+$
$\Xi_c'^0$	$\Xi_c^0 \gamma$	2578.8 ± 0.5		$\frac{1}{2}^+$
$\Xi_c(2645)^+$	$\Xi_c^0 \pi^+$	2645.53 ± 0.31	2.14 ± 0.19	$\frac{3}{2}^+$
$\Xi_c(2645)^0$	$\Xi_c^+ \pi^-$	2646.32 ± 0.31	2.35 ± 0.22	$\frac{3}{2}^+$
$\Xi_c(2790)^+$	$\Xi_c'^0 \pi^+$	2792.0 ± 0.5	8.9 ± 1.0	$\frac{1}{2}^-$
$\Xi_c(2790)^0$	$\Xi_c'^+ \pi^-$	2792.8 ± 1.2	10.0 ± 1.1	$\frac{1}{2}^-$
$\Xi_c(2815)^+$	$\Xi_c^+ \pi^+ \pi^-$, $\Xi_c(2645)^0 \pi^+$, $\Xi_c^{0'} \pi^+$	2816.67 ± 0.31	2.43 ± 0.26	$\frac{3}{2}^-$
$\Xi_c(2815)^0$	$\Xi_c^0 \pi^+ \pi^-$, $\Xi_c(2645)^+ \pi^-$, $\Xi_c^{+'} \pi^-$	2820.22 ± 0.32	2.54 ± 0.25	$\frac{3}{2}^-$
$\Xi_c(2930)^0$	$\Lambda_c^+ K^-$	$2928.9^{+3.1}_{-12.4}$	19.5^{+10}_{-12}	
$\Xi_c(2970)^+$	$\Lambda_c^+ K^- \pi^+$, $\Sigma_c^{++} K^-$, $\Xi_c(2645)^0 \pi^+$, $\Xi_c^{0'} \pi^+$	2969.4 ± 0.8	$20.9^{+2.4}_{-3.5}$	
$\Xi_c(2970)^0$	$\Xi_c(2645)^+ \pi^-$, $\Xi_c^{+'} \pi^-$	2967.8 ± 0.8	$28.1^{+3.4}_{-4.0}$	
$\Xi_c(3055)^+$	$\Sigma_c^{++} K^-$, ΛD^+	3055.9 ± 0.4	7.8 ± 1.9	
$\Xi_c(3055)^0$	ΛD^0	3059.0 ± 0.8	6.4 ± 2.4	
$\Xi_c(3080)^+$	$\Lambda_c^+ K^- \pi^+$, $\Sigma_c^{++} K^-$, $\Sigma_c(2520)^{++} K^-$, ΛD^+	3077.2 ± 0.4	3.6 ± 1.1	
$\Xi_c(3080)^0$	$\Lambda_c^+ K_S^0 \pi^-$, $\Sigma_c^0 K_S^0$, $\Sigma_c(2520)^0 K_S^0$	3079.9 ± 1.4	5.6 ± 2.2	


Ω_c Family



$$\Omega_c^{*0} \rightarrow \Omega_c^0 \gamma$$

$[70.8 \pm 1.0(\text{stat.}) \pm 1.1(\text{syst.})] \text{ MeV}/c^2$



[B. Aubert *et al.* (BaBar Collaboration),
Phys. Rev. Lett. **97**, 232001 (2006)]



$$\Delta M_{\Omega_c^0} = [70.7 \pm 0.9(\text{stat.})^{+0.1}_{-0.9}(\text{syst.})] \text{ MeV}/c^2$$

[E. Solovieva, R. Chistov *et al.* (Belle
Collaboration), Phys. Lett. B **672**, 1 (2009)]

Ω_c Family



Resonance	Mass (MeV)	Γ (MeV)	N_σ
$\Omega_c(3000)^0$	$3000.4 \pm 0.2 \pm 0.1^{+0.3}_{-0.5}$	$4.5 \pm 0.6 \pm 0.3$	20.4
$\Omega_c(3050)^0$	$3050.2 \pm 0.1 \pm 0.1^{+0.3}_{-0.5}$	$0.8 \pm 0.2 \pm 0.1$	20.4
		< 1.2 MeV, 95% CL	
$\Omega_c(3066)^0$	$3065.6 \pm 0.1 \pm 0.3^{+0.3}_{-0.5}$	$3.5 \pm 0.4 \pm 0.2$	23.9
$\Omega_c(3090)^0$	$3090.2 \pm 0.3 \pm 0.5^{+0.3}_{-0.5}$	$8.7 \pm 1.0 \pm 0.8$	21.1
$\Omega_c(3119)^0$	$3119.1 \pm 0.3 \pm 0.9^{+0.3}_{-0.5}$	$1.1 \pm 0.8 \pm 0.4$	10.4
		< 2.6 MeV, 95% CL	
$\Omega_c(3188)^0$	$3188 \pm 5 \pm 13$	$60 \pm 15 \pm 11$	


[R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. Lett. **118**, 182001 (2017)]

Ω_c Family

$$\Omega_c(X)^0 \rightarrow \Xi_c^+ K^-$$

↓


$$\Xi^-\pi^+\pi^+, \Lambda K^-\pi^+\pi^+, \Xi^0\pi^+, \Xi^0\pi^+\pi^-\pi^+, \Sigma^+K^-\pi^+, \Lambda K_S^0\pi^+, \Sigma^0K_S^0\pi^+$$



Ω_c Excited state	3000	3050	3066	3090	3119	3188
Yield	37.7 ± 11.0	28.2 ± 7.7	81.7 ± 13.9	86.6 ± 17.4	3.6 ± 6.9	135.2 ± 43.0
Significance	3.9σ	4.6σ	7.2σ	5.7σ	0.4σ	2.4σ
LHCb mass	$3000.4 \pm 0.2 \pm 0.1$	$3050.2 \pm 0.1 \pm 0.1$	$3065.5 \pm 0.1 \pm 0.3$	$3090.2 \pm 0.3 \pm 0.5$	$3119 \pm 0.3 \pm 0.9$	$3188 \pm 5 \pm 13$
Belle mass (with fixed Γ)	$3000.7 \pm 1.0 \pm 0.2$	$3050.2 \pm 0.4 \pm 0.2$	$3064.9 \pm 0.6 \pm 0.2$	$3089.3 \pm 1.2 \pm 0.2$...	$3199 \pm 9 \pm 4$

Ξ_{cc} Family


$$\Xi_{cc}^+ \rightarrow \Lambda_c^+ K^- \pi^+$$



[M. Mattson *et al.* (SELEX Collaboration), Phys. Rev. Lett. **89**, 112001 (2002)]

$$m_{\Xi_{cc}^+} = (3518.7 \pm 1.7) \text{ MeV}/c^2$$


$$\Xi_{cc}^+ \rightarrow p D^+ K^-$$



[A. Ocherashvili *et al.* (SELEX Collaboration), Phys. Lett. B **628**, 18 (2005)]

Ξ_{cc} Family

$$\Xi_{cc}^{++} \rightarrow \Lambda_c^+ K^- \pi^+ \pi^+$$



$$m_{\Xi_{cc}^{++}} = (3621.40 \pm 0.72[\text{stat.}] \pm 0.27[\text{syst.}] \pm 0.72[\Lambda_c^+]) \text{ MeV}/c^2$$

$$m_{\Xi_{cc}^{++}} - m_{\Xi_{cc}^+} = (103 \pm 2) \text{ MeV}/c^2$$

[R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. Lett. **119**, 112001 (2017)]

Conclusions

- The Ξ_{cc} state reported by LHCb is consistent with most theoretical expectations, but it is inconsistent with being an isospin partner to the Ξ_{cc} state reported previously by the SELEX Collaboration.
- Recently observed excited Ω_c states present a unique opportunity to test and further improve theoretical models, that predict properties of heavy hadrons.
- More accurate Ξ_c mass values is of both practical and theoretical interest, and knowing their widths can then lead to measurements of the matrix elements of their decays. These matrix elements are also applicable to other excited charm and bottom baryons.
- No direct measurements of the J^P of any of the excited strange charmed baryons are available. Constraints on the quantum numbers can be inferred only from the decay pattern.
- Interesting feature is that highly excited charmed baryons can decay to a charm meson and a non-charm baryon.